

GUIDANCE AT A GLANCE - *CLOSTRIDIUM DIFFICILE*

These guidelines support the control and prevention of *C.difficile* in community and primary care settings.

Key Ref: PHE (2013) Updated guidance on the management and treatment of *Clostridium difficile* infection

KEY POINTS

- Early diagnosis prevents complications and saves lives
- Prudent prescribing of antibiotics may prevent infection
- Communication of infection risk helps prevent cross infection
- Alcohol hand rub is **ineffective** at killing *C.difficile* spores.
- Risk factors for infection = over 65s, recent hospitalisation, recent antibiotics, GI procedures and gastric ulcer medications

These affect the balance of bacteria in the bowel providing an opportunity for C.difficile to multiply, produce toxin and inflame the bowel

ANTIBIOTICS

My patient has diarrhoea

A useful mnemonic protocol for potentially infective diarrhoea):
Suspect that a cause may be infective where there is no clear alternative cause of diarrhoea or the patient has recently received antibiotics
Isolate the patient if appropriate – *i.e. care home residents*
Gloves and aprons should be worn to reduce cross contamination
Hand washing with soap & water before and after each patient contact and the patient's environment
Test the stool for toxin, by sending a specimen immediately

Watery or mucoid diarrhoea with or without blood (typical smell and green appearance), abdominal pain, loss of appetite, fever.

CDI Symptoms

My patient has confirmed CDifficile Infection

- **No repeat specimens** are required once diagnosed. For **toxin gene detected** results, treat as CDifficile infection (CDI) if symptomatic.
- Review the need for any current antibiotics and stop the course if possible – if unable to stop, change to a narrow spectrum antibiotic.
- Review other drugs that may potentially cause diarrhoea.
- Proton pump inhibitors (PPI) should be reviewed/reduced where possible
- For treatment options refer to the algorithm over and also the local antimicrobial guidelines for primary care.
- Maintain hydration, monitor diarrhoea (*for care homes residents there is a care plan, fluid balance chart and Bristol Stool Chart to support this*)
- Where patients/carers are unable to manage due to the symptoms of *C.difficile* consider referral to social services.

Communicating infection risk:

- If admitting a patient with symptoms, notify the receiving area so appropriate isolation can be instigated. Use the transfer sticker.
- The IPC team will write to your patient supplying a leaflet and card – copied to the practice.
- Record the infection risk on the patient record as guided by the IPC team

CDI can lead to: dehydration, electrolyte imbalance low blood albumin pseudomembranous colitis, toxic megacolon, sepsis, death.

Complications


My patient has a history of CDifficile and needs antibiotics

There is an increased risk of CDI if someone has had it before or has had the gene detected. Consider narrow spectrum antibiotics if treating other infections

Resources

- Antimicrobial guidelines
- Patient held card
- CDifficile patient information leaflet
- Transfer stickers
- PHE guidelines '13

Treatment Algorithm for CDI



Severity

Mild CDI is not associated with a raised WCC; it is typically associated with <3 stools of types 5-7 on the Bristol Stool Chart per day.

Moderate CDI is associated with a raised WCC that is $<15 \times 10^9/L$; it is typically associated with 3-5 stools per day.

Severe CDI is associated with a WCC $>15 \times 10^9/L$, or an acute rising serum creatinine (i.e. >50% increase above baseline), or a temperature $>38.5^\circ C$, or evidence of colitis (abdominal or radiological signs). The number of stools may be a less reliable indicator of severity.

Life-threatening CDI includes hypotension, partial or complete ileus or toxic megacolon, or CT evidence of severe disease.

N.B. Mild/moderate CDI – treat as non-severe.